References :

1. Abdel-Mohsen H.T., Ragab F.A.F., Ramla M.M, Diwani H.I.E. (2010)., synthesised novel 2, 4, 5- substituted pyrimidine derivatives. 2, 4, 5-substituted pyrimidine., *Eur J Med Chem*; **45,** 2336–2344.
2. Abdullah E.S.A., (2011) Synthesized 2-thioxopyrimidine derivatives, *Molecule*; **16,** 3416-3419.
3. Ahmed O.M, Mohamad M.A.,(2009) reported the synthesis of (1, 3, 5-tetrahydro-4, 1-benzoxazepine-3-yl)-pyrimidines and evaluated for anticancer activity, *Eur Med Chem*.; **44,** 3519-3523.
4. Ajam M.A, Kumar B.P.R, Suresh B.,(2008) Synthesised pyrido [2, 3-d] pyrimidines., *Ind J Pharm Sci*.; 672-677.
5. Amin KM, El-Zahar M.I, Anwar M.M, Kamel M.M, Mohamed M.H.,(2009) Synthesis, antitumor activity and molecular docking study of novel benzofuran-2-yl pyrazole pyrimidine derivatives., *Acta Poloniae Pharmaceutica and Drug Research.*; **66,** 279 -291.
6. Amr A.G.E, Mohamed A.M, Mohamed S.F, Hafez NAA, Hammam AG.,(2006) Synthesised pyrimidine bridged thiadiazole derivatives .5-{(4, 6-disubstituted pyrimidine-2-yl) thio methyl}-Nphenyl-1, 3, 4, thiadiazol-2-amines *Bioorg Med Chem*.; **14,** 5481–5488.
7. Atwal, K. S.; O'Reilly, B. C.; Gougoutas, J. Z.; Malley, M. F.,(1987) Very attractive approach to the synthesis of Biginelli compounds was developed by Atwal and co-workers., *Heterocycles*.;**26,** 1189- 1192.
8. Atwal, K. S.; Swanson, B. N.; Unger, S. E.; Floyd, D. M.; Moreland, S.; Hedberg, A.; O'Reilly, B. C.,(1991),5-acyl-1,2,3,4-tetrahydropyrimidine-2-thiones/ones and these compounds are very potent calcium channel blockers., *J.Med. Chem.*; **341**,806.
9. Becerro, M. A.; Uriz, M. J.; Turon, X. 1994,seasonal patterns of toxicity in benthic invertibrates :the encrusting sponge crambe crambe (poecilosclerida) *Mar. Biol.*, **121,** 301-307
10. Berlinck, R. G. S.,(2002), Among the most notable of these are the crambescidin., *Nat. Prod. Rep.*; **19,** 617
11. Biginelli, P.,(1893), Synthesis of 5-acyl-1,2,3,4-tetrahydropyrimidine-2-thiones/ones., *Gazz. Chim. Ital.,* **23,** 360-416.

12. Cho, H.; Ueda, M.; Shima, K.; Mizuno, A.; Hayashimatsu, M.; Ohnaka, Y.;Hamaguchi, M.; Aisaka, K.; Hidaka, T.; Kawai, M.; Takeda, M.; Ishihara, T.;Funahashi, K.; Satoh, F; Morita, M.; Noguchi, T., (1989) Ring cleavage of dihydropyrimidine skeleton., *J.Med. Chem*., **32**, 2399.

13. (a) Coffey, D. S.; McDonald, A. I.; Overman, L. E.; Rabinowitz, M. H.;Renhowe, P A. J. (2000) *Am. Chem. Soc.*(2000), **122,** 4893. (b) Coffey, D. S.; Overman, L. E.; Stappenbeck, F. J. *Am. Chem. Soc.,* **122,** 4094.

14. Ding S, Gray NS, Ding Q, Schultz PG. (2001) Expanding the diversity of purine libraries. *Tetrahedron Lett*; **42**, 8751-5

15. (a) Drews, *J. Science* (2000), **287**, 1960; (b) Wess, G.; Urmann M.; Sickenberger, B. Angew. Medicinal Chemistry: Challenges and Opportunities., *Chem. Int. Ed*. (2001), 40, 3341.

16. Evans, B. E.; Rittle, K. E.; Bock, M. G.; Dipardo, R. M.; Freidinger, R. M.; Whitter, W. L.; Lundell, G. F.; Veber, D. F.; Anderson, P. S.; Chang, R. S. L.; Lotti, V. J.; Cerino, D. J.; Chen, T. B.; Kling, P. J.; Kunkel, K. A.; Springer, J. P.; Hirshfield, J., (1988) describe selected structural types that bind to multiple, unrelated classes of protein receptors and enzymes as high affinity ligands *J. Med. Chem*., **31,** 2235.

17. Folkers, K.; Johnson, T. B. (1933) the primary bimolecular condensation product of benzaldehyde and urea is the first intermediate in this reaction. *J. Am. Chem. Soc*.**, 55**, 3781-3791.

18. Folkers, K.; Johnson, T. B, (1933) the primary bimolecular condensation product of benzaldehyde and urea is the first intermediate in this reaction. *J. Am. Chem.Soc.*, **55,** 3781-3791

19. Gangjee A ,Kurup S, Ihnat M.A, Thorpe J.E, Shenoy S.S. (2010), synthesis of Pyrazolo [1,5-a] pyrimidine derivative., *Bioorg Med Chem,*; **18,** 3575–3587.

20. Ghorab M.M, Ragab F.A, Alqasoumi S.I, Alafeefy A.M, Aboulmagd S.A. (2010), Synthesis of Some New Pyrazolo[3,4-d]pyrimidine., *Eur J Med Chem,*; **45,** 171–178.

21. Hashem A.A, Youssef Hoda M.M, Hussein A.R (2011)., synthesis of some new thiazolopyrimidine, Pyrrolothiazolo pyrimidine & thiazlopyrrolothiazolopyrimidine derivatives.. *J Chinese Chem Soc*,, **58:** 41-48.

22. Horton, D. A.; Bourne, G. T.; Smythe, M. L. (2003), benzothiophenes., *Chem. Rev.*, **103,** 893.

23. Hu, E. H.; Sidler, D. R.; Dolling, U. H.(1998), substituted aromatic or aliphatic aldehydes and thiourea were employed *J. Org. Chem*., **63,** 3454-3457.
24. Hurst, E. W.; Hull, R. (1961) An expedient protocol of the Biginelli dihydropyrimidine synthesis, .*J. Med. Pharm. Chem*., **3,** 215.
25. Kappe, C. O. (1993) 100 Years of of the biginelli dihydropyrimidine synthesis, *Tetrahedron*.
26. Kappe, C. O.; Fabian, M. F. (1997), Conformational analysis of 4-aryl-dihydropyrimidine calcium channel modulators. A comparison of ab initio, semiempirical and X-ray crystallographic studies, *Tetrahedron, ,* **53(8),** 2803.
27. Kappe, C. O.; Falsone, F.; Fabian, W. M. F.; Belaj, F.(1999) Isolation, Conformational Analysis and X-Ray Structure Determination of a Trifluoromethylstabilized Hexahydropyrimidine-An Intermediate in the Biginelli Reaction.. *Heterocycles*, **51**,77-84.
28. (a) Kashman, Y.; Hirsh, S.; McConnell, O. J.; Ohtani, I.; Kusumi, T.;Kakisawa, H. (1989),synthesis of ptilomycalin A., *J. Am. Chem. Soc.*, 111,8925. (b) Jares-Erijman, E. A.;Sakai, R.; Rinehart, K. L(1991). *J. Org. Chem*.**, 56,** 5712.
29. Kolbn H. C.; Finn, M. G.; Sharpless, K. B. Angew. (2001), diverse chemical function from a few good reactions, *Chem. Int. Ed.*, **40,** 2004.
30. Kraljevic TG, Krištafor S, Šuman L, Kralj M, Ametamey SM, Cetina M, Malic SR. (2010), 5, 6-disubstituted pyrimidine derivative were synthesized, *Bioorg Med Chem,* **18**, 2704–2712.
31. Lombardino, J. G.; Lowe III, (2004), *J. A. Nat. Rev*. *Drug Discov.*, **3**, 853.
32. Mohamed N.R, El-Saidi MMT, Alia YM, Elnagdi MH. (2007), the synthesis of N-4-phenyl substituted-6(2,4-dichlorophenyl methyl)-7H-Pyrrolo[2,3-d]pyrimidine-2,4-diamines and evaluated for anticancer activity. *Bioorg Med Chem,*; **15,** 6227–6235.
33. (a) Nagasawa, K.; Georgeiva, A.; Koshino, H.; Nakata, T.; Kita, T.;Hashimoto, Y. (2002) Concise Synthesis of α-Trisubstituted Amines from Ketones Using *N*-Methoxyamines *Org. Lett.*, **4**, 177.(b) Ishiwata, T.; Hino, T.; Koshino, H.;Hashimoto, Y.; Nakata, T.; Nagasawa, K. (2002) *Org. Lett*., **4**, 2921. (c) Nagasawa, K.; Ishiwata, Y.; Hasimoto, Y.; Nakata, T. (2002) *Tetrahedron Lett.*, **43,** 6383. (d) Moore, C. G.; Murphy, P. J.; Williams, H. L.; McGown, A. T.; Smith, N. K. (2003) *Tetrahedron Lett.***, 44,** 251.
34. Nicolaou, K. C.; Pfefferkorn J. A.; Roecker, A. J.; Cao, G. Q.; Barluenga, S.;

Mitchell, H. J. (2000), Synthesis of various drug like substituted benzo pyrans. *J. Am. Chem. Soc.,* **122,** 9939.

35. Overman, L. E.; Rabinowitz, M. H.; Renhowe, P. A, (1995) An expedient protocol of the Biginelli dihydropyrimidine synthesis using carbonyl equivalents., *J. Am. Chem. Soc.,* , **117,** 2675

36. Patil, A. D.; Kumar, N. V.; Kokke, W. C.; Bean, M. F.; Freyer, A. J.; De Brosse, C.; Mai, S.; Truneh, A.; Faulkner, D. J.; Carte, B.; Breen, A. L.;Hertzberg, R. P. Johnson, R. K.; Westley, J. W.; Potts, B. C. M. (1995), the 2-oxodihydropyrimidine-5-carboxylate core unit is found in nature and in potent HIVgp-120-CD4 inhibitors *J.Org. Chem.*, **60**, 1182.

37. (a) Patil, A. D.; Kumar, N. V.; Kokke, W. C.; Bean, M. F.; Freyer, A. J.; De Brosse, C.; Mai, S.; Truneh, A.; Faulkner, D. J.; Carte, B.; Breen, A. L.;Hertzberg, R. P. Johnson, R. K.; Westley, J. W.; Potts, B. C. M. (1995), the 2-oxodihydropyrimidine-5-carboxylate core unit is found in nature and in potent HIVgp-120-CD4 inhibitors *J.Org. Chem.*, **60**, 1182. (b) Patil, A. D.; Freyer, A. J.; Taylor, P. B.; Carte, B.; Zuber,G.; Johnson, R. K.; Faulkner, D. J., batzelladine. (1997), *J. Org. Chem*., **62,** 1814.

38. Ramesh B, Kulakani SV, Ramachandra R. (2010), synthesis of novel benzimidazole pyrimidine conjugates as potent antitumor agent., *Int J Pharma Sci*,**; 2(1),** 426-428.

39. Rovnyak, G. C.; Atwal, K. S.; Hedberg, A.; Kimball, S. D.; Moreland, S.;Gougoutas, J. Z.; O'Reilly, B. C.;Schwartz, J.; Malley, M. (1981), Ring cleavage of dihydropyrimidine skeleton., F. *J. Med. Chem*., **46,** 3433.

40. Shutalev, A. D.; Kishko, E. A.; Sivova, N. V.; Kuznetsov, A. Yu. (1998), A New Convenient Synthesis of 5-Acyl-1,2,3,4-tetrahydropyrimidine-2-thiones/ones., *Molecules*, **3,** 100-106.

41. Sinder, B. B.; Shi, Z. (1993), "Biomimetic Synthesis of (±)-Crambines A, B, C1, and C2. Revision of the Structure of Crambines B and C1", *J Org. Chem.,* , **58,** 3828

42. Sweet, F.; Fissekis, J. D. (1973) Fissekis' Mechanism, Evidence against Reaction with thiourea *J. Am. Chem. Soc.,* **95,** 8741.